Cephora

Cephora is a maincrop variety with a very good taste, French fry quality and excellent presentation. The tubers are oval to long-oval. The variety has very good resistance to late blight.





Q Impala 95T118-2 ð



potato driven

Basic properties		
Dormancy	6	
Foliage development	6	
Maturity	6	
Tuber shape	OLO	
Uniformity of tubers	7	
Tuber size	7	
Uniformity of size	7	
Relative tubercount	7	
Shallowness of eyes	7	
Colour of skin	Y	
Colour of flesh	LY	
Early yield %	115	
Mature yield %	100	

PCN Rostochiensis	1-
PCN Pallida	-
Wart disease	-
Late blight foliage	8
Late blight tuber	5,
PVY	7,
Common scab	5
Fusarium	7
Tobacco Rattle Virus	-
Mechanical damage	7
Second growth	7
Bruising	6

Cultivation and storage (point of attention)

Seed treatment Plot selection
Fertilisation
Planting
Cultivation
Metribuzin
Harvesting
Storage

Point of Attention

Handling

Pre-sprouting not required Avoid soils with high common scab potential

N±/ P+/K++ Average planting distance, but plant deeper (4-5 cm) Well shaped ridges with good soil coverage are required Average sensitive

Quick drying after harvesting is important No special handling required

Quality

S	kin presentation	7,5
U	WW	375
D	ry matter %	20,5
С	ooking type	AB/B
Q	uality of taste	7,5
С	risps index	5
F	rench fries index	7

Explanation

General:	9 (=favorable) 1 (=unfavorable)
Tuber shape:	R (=round) O (=oval) L (=long)
Colour:	D (=dark) Y (=yellow) L (=light) C (=cream) W (=white) R (=red) P (=purple) Br (=brown)

Segment



Plantera B.V. does not accept liability for any loss or damages caused by the use of this data.